



Advancing Transfusion and
Cellular Therapies Worldwide

Association Bulletin #17-02

Date: March 16, 2017
UPDATED MARCH 2026

To: AABB Members

From: Zbigniew Szczepiorkowski, MD, PhD – President
Thomas Hopkins, MBA – Interim Chief Executive Officer

Re: Updated Strategies to Limit or Prevent Iron Deficiency in Blood Donors

This bulletin, developed by the AABB Donor Health and Safety Committee and approved by the AABB Board of Directors, contains information for all establishments that collect Whole Blood (WB) or apheresis component donations. It supersedes Association Bulletin #12-03 “Strategies to Monitor, Limit, or Prevent Iron Deficiency in Blood Donors” and recommends action(s) to limit or help prevent iron deficiency in blood donors.

Summary

Association Bulletin #12-03 recommended that blood collection establishments take voluntary action(s) to monitor, limit, or prevent iron deficiency in blood donors. AABB’s current version of the *Standards for Blood Banks and Transfusion Services* require procedures to ensure donors are given, and acknowledge reading, educational materials that include the risks of postdonation iron deficiency.¹ Mounting evidence indicates that frequent donors require iron supplementation to prevent significant iron depletion that may have deleterious health consequences. Blood collection establishments should therefore take action(s) with all donors or selected subpopulations at risk for, or from, iron deficiency (listed below) to include at least one of the following interventions/strategies: 1) development of programs to provide replacement iron in the absence of ferritin measurements, 2) evidence-based lengthening of the interdonation interval and/or restriction of the number of donations per year, or 3) measurement of serum or plasma ferritin leading to recommended actions (including iron supplementation) for donors with low ferritin levels.



Advancing Transfusion and
Cellular Therapies Worldwide

Association Bulletins, which are approved by the AABB Board of Directors, may include announcements of standards or requirements for accreditation, recommendations on emerging trends or best practices, and/or pertinent information. This bulletin contains information and recommendations.

Background

Iron deficiency is observed in blood donors, with rates ranging from 49 percent in men to 66 percent in women, who donate frequently.^{2,3} Premenopausal women are at particular risk for developing iron deficiency. As iron deficiency progresses, it eventually results in anemia, but the potential for adverse health consequences exists before anemia occurs.^{4,5} Iron deficiency (particularly in the presence of anemia) can affect several physiological and health parameters. Adverse effects include: cognitive dysfunction, fatigue, pregnancy-related complications (perinatal mortality, preterm delivery, low birthweight, newborn cognitive abnormalities), decreased exercise endurance, pica (the desire to compulsively ingest non-food substances such as ice or clay), and restless legs syndrome.⁶⁻²³

Iron deficiency in blood donors exists along a continuum, ranging from reduced iron stores to iron-deficiency anemia. The likelihood and severity of iron depletion are influenced by donation frequency, sex, age, baseline iron status, and cumulative red cell loss. Recognition of this graded risk is important for implementing proportionate and effective donor protection strategies.²

Subgroups at Risk

The following donors are at risk for developing iron deficiency and/or at increased risk for adverse effects of iron deficiency:

- Young donors.
- Premenopausal females.
- Frequent donors.
- Donors with hemoglobin values near the minimum for eligibility.

Pertinent Studies

Many of the studies reporting adverse effects are more suggestive than definitive, particularly in non-anemic iron deficiency (NAID).⁴ Most iron-deficient blood donors have NAID, and a growing body of literature identifies the effect of NAID in impaired quality-of-life



Advancing Transfusion and
Cellular Therapies Worldwide

measures.^{9,19,23} Frank iron deficiency anemia occurs occasionally in blood donors and is more clearly and consistently associated with adverse outcomes. Although more randomized, controlled trials are needed, our current understanding of the potential harms associated with NAID resulting from blood donation demands thoughtful clinical and ethical attention.²⁴

Since Association Bulletin #12-03 was published, two important randomized, controlled trials investigating iron balance and iron recovery in frequent blood donors have been published. The first study, conducted as part of the National Heart, Lung, and Blood Institute (NHLBI)-supported Recipient Epidemiology and Donor Evaluation Study-III (REDS-III) was the HEmoglobin Iron and Recovery Study (HEIRS). The second study, also funded by the NHLBI, was the Strategies To Reduce Iron DEficiency (STRIDE) study.^{25, 26} Preliminary data from another NHLBI-funded study, the Comparison of the History of Donation and Iron LeveLs in Teen Blood Donors (CHILL) study, have also been reported.²⁷

- HEIRS

Important aims of HEIRS were to determine the time to postdonation hemoglobin and ferritin recovery with and without oral iron supplementation.²⁵ HEIRS enrolled 136 female and 79 male blood donors ≥ 18 years old at four US blood centers. None of the donors had donated blood in the prior 4 months. Donors were separated into two groups by storage iron level at enrollment: a lower ferritin group (≤ 26 ng/mL; iron-depleted) and a higher ferritin group (> 26 ng/mL; not iron-depleted). At their next study visit (3-8 days after donation), subjects were randomly assigned to receive either 37.5 mg of elemental iron as oral ferrous gluconate daily for 24 weeks, or no supplementation within strata based upon initial ferritin level, gender, and age. Hemoglobin and ferritin levels were measured seven times during the study, with the final measurement 24 weeks after donation. Approximately 90 percent of subjects completed at least six of the seven scheduled visits.

Primary results of the study are as follows. Compared to those who did not take iron, donors taking supplements returned to 80 percent of predonation hemoglobin levels faster in both the lower ferritin group (median of 5 weeks vs 23 weeks) and higher ferritin group (median of 4 weeks vs 11 weeks). Similarly, donors taking iron supplements recovered lost iron (as determined by ferritin measurements) more rapidly than those not receiving supplements (11 weeks vs > 24 weeks for return to baseline ferritin). At 24 weeks after donation, two-thirds of the donors who did not take supplements still had not recovered the iron lost at donation. Thus, in addition to showing the efficacy of iron supplementation, HEIRS demonstrated that extending the interdonation interval to 12 or



Advancing Transfusion and
Cellular Therapies Worldwide

16 weeks is not sufficient for the average donor to recover iron stores without supplementation.

In a second analysis, the investigators calculated the total body iron, the sum of storage iron and iron in hemoglobin.²⁸ They identified a statistically significant effect of iron supplementation on total body iron recovery in the first 8 of the 24 weeks of iron administration. This effect was demonstrable (but weakly) in the second 8 weeks only in those with iron deficiency, and not in any donors by the third 8-week period. Taken together, these findings suggest that the majority of iron's beneficial effect is achieved within the first 8 weeks of supplementation. Thus, daily intake of low-dose oral iron for 8 weeks is likely to be an effective way to remedy donation-related iron depletion.

- STRIDE

The STRIDE study was a multicenter, randomized, blinded, and placebo-controlled study investigating several readily implemented approaches to mitigate donor iron deficiency.²⁶ Frequent blood donors from three blood centers were randomly assigned to one of five study arms. Donors in two intervention arms received either 19 mg or 38 mg of elemental iron as ferrous gluconate for 60 days after each donation. The third intervention arm was informational; in this arm (designated as the "Iron Status Information" arm), donors received a letter informing them of their ferritin level. The letter recommended continued frequent donation for donors with ferritin levels ≥ 26 ng/mL, while donors with ferritin levels < 26 ng/mL were advised to take self-purchased iron supplements and/or delay donation for 6 months. In the two control arms, donors received a letter encouraging frequent donation or administration of a placebo in lieu of iron supplements. Donors were evaluated at multiple follow-up visits over a 2-year period. Ferritin, soluble transferrin receptor, and a complete blood count were measured at each donation. Of 692 enrolled donors, 57 percent completed the study. There was a higher dropout rate in the three groups receiving pills, including the group taking placebo, unrelated to side effects that occurred equally in all three groups.

Ferritin level and hemoglobin concentration at the end of the study were statistically equivalent among subjects randomly assigned to the 19- and 38-mg iron groups and the Iron Status Information arm. Values in the placebo and no-information-letter study arms were significantly lower. This finding held true for other iron deficiency measurement cutoffs. Reductions of 30-80 percent in the prevalence of iron deficiency from enrollment occurred only in the iron and Iron Status Information study arms. Additional analyses of absolute changes in iron and hemoglobin status from enrollment to final visit and models



Advancing Transfusion and
Cellular Therapies Worldwide

incorporating data from all visits by all enrolled subjects (not just those completing the study) showed that donors in the Iron Status Information group did not improve their iron status and hemoglobin concentration quite as much as those who were directly provided iron pills. Differences between the primary and secondary analyses can be attributed to the diverse responses of Iron Status Information donors, some of whom took iron and/or lengthened their interdonation intervals, while others did neither, particularly those with ferritin levels ≥ 26 ng/mL. Taken as a whole, STRIDE results demonstrated that providing 19 or 38 mg of daily iron or iron status information after ferritin testing was effective and mostly equivalent in mitigating postdonation iron deficiency.

- CHILL

Recently, the REDS-III program completed the CHILL study.²⁷ Two blood centers enrolled donors at high school blood drives during 2015-2016 and measured donor ferritin levels at their first and any subsequent donations during the school year. In total, 4,265 donors making 6,219 donations were enrolled. Of those, 3,714 donors making 5,439 donations were aged 16-18 (defined as teen donors) while the remainder were adult donors, aged 19-49, who served as controls.

Preliminary data analysis indicated that at study outset, the mean ferritin levels of first-time donors (at each age and gender) did not differ from national population data obtained in the 2001-2002 National Health and Nutrition Examination Survey (NHANES) but that the prevalence of iron deficiency was greater in teen donors than in the adult control donors. Further, with repeat donation, the prevalence of iron deficiency was also greater in female and male teen donors than in gender-matched adults. CHILL investigators concluded that the baseline prevalence of iron deficiency is higher in teenagers before donation and may be exacerbated to a greater extent by blood donation when compared to adult donors; this latter preliminary finding is undergoing further evaluation by statistical modeling.

Post-2017 Evidence and National Implementation Experience

The REDS-II and REDS-III studies (RISE, HEIRS, STRIDE, CHILL) established that iron deficiency is common in blood donors, often precedes anemia, and can be mitigated by iron supplementation or donation-interval modification. Since those studies, several blood services have moved from trial environments to national implementation of ferritin-based donor management, generating high-quality real-world data.



Advancing Transfusion and
Cellular Therapies Worldwide

These programs demonstrate that ferritin testing and ferritin-guided interventions are feasible in routine practice and are associated with meaningful reductions in iron deficiency prevalence among donors. More recent randomized controlled trials and program evaluations conducted outside the United States confirm that ferritin-guided approaches, including donation-interval modification and iron supplementation, effectively mitigate donation-related iron depletion while maintaining donor participation. Importantly, these studies extend evaluation beyond laboratory recovery to include outcomes such as hemoglobin deferral rates, donor return patterns, and donor wellbeing.

Key developments include:

- National ferritin-based policies implemented at scale demonstrate feasibility, donor acceptance, and meaningful reductions in iron deficiency prevalence.
- Randomized controlled trials evaluating ferritin-guided donation intervals and iron supplementation strategies under operational conditions confirm that ferritin-based approaches effectively protect donors while preserving donation activity.
- System-level outcomes, including reduced low-hemoglobin deferrals and improved donor return patterns, have now been documented, extending the focus beyond laboratory recovery alone.

Examples of key studies:

1. Large national implementation data from the Netherlands (Sanquin) show that ferritin-based policies are feasible at scale and materially change donor management. In their ferritin-based deferral policy, ferritin is measured (notably including first-time donors and periodic repeat testing), and donors with low ferritin are deferred for months depending on severity; early results describe program effects and the distribution of low ferritin in donors under an operational policy rather than a trial setting.²⁹
2. The Netherlands also progressed from policy to randomized evaluation: ferritin-guided strategies, including supplementation approaches, have been formally trialed, including a double-blind RCT reported in *The Lancet Haematology* in 2025.³⁰⁻³¹
3. Denmark provides complementary “system-level” evidence: ferritin-guided supplementation introduced in routine practice for female first-time donors reduced subsequent low-Hb occurrence and was associated with improved donation activity, which is important because it links donor protection to operational outcomes (deferrals and supply).³²
4. A 2024 comparative European overview of donor-iron management shows that countries are now using materially different strategies (supplementation vs ferritin-guided interval



Advancing Transfusion and
Cellular Therapies Worldwide

policies), and it helps position the AABB approach in an international context, including the consequences of not using either ferritin-guidance or supplementation (higher deferral patterns).³³

5. The dosing science for oral iron has matured: alternate-day dosing improves absorption in iron-depleted women compared with consecutive-day dosing, consistent with hepcidin physiology. This matters for donor programs because it offers a way to reduce GI side effects while maintaining repletion efficiency, which can improve adherence in donors.³⁴

Recommendations

Ethical Framework

Preventing donation-related iron deficiency should be viewed as a donor-safety obligation, comparable to other established donor-protection measures. Protecting donors from foreseeable, preventable harm supports donor trust, program sustainability, and the ethical foundation of voluntary blood donation.

In light of 1) the possibility that some donors may develop iron deficiency anemia after donation, 2) the known progression of iron deficiency that occurs with ongoing frequent blood donation, and 3) the potential for adverse effects of iron deficiency, blood collection establishments should develop policies to limit or help prevent iron deficiency in their donors. These policies should go beyond the current AABB standard requiring development of educational materials for all donors on the risk of postdonation iron deficiency.

Recommendation #1: Blood Collection Establishments Should Prepare Comprehensive Educational Materials to Assist Donors in Better Understanding the Need for Specific Interventions.

Blood collection establishments should provide all donors with comprehensive information that not only includes information on the risks of iron deficiency resulting from blood donation and their applicability to particular at-risk donor subgroups (as described below), but also discusses the benefits of iron supplementation following donation. This information should include recommendations on the type of iron, dosage, and length of supplementation required to replace an equivalent amount of iron lost at donation. Because iron supplements may be harmful in some individuals or mask conditions associated with gastrointestinal blood loss, donors with a personal or



Advancing Transfusion and
Cellular Therapies Worldwide

family history of hereditary hemochromatosis, familial polyposis, or colorectal cancer should be advised to check with their health-care provider before commencing iron supplementation. Donors should also be advised to consult with their health-care provider or pharmacist about the effect of iron on absorption of other medications.

Recommendation #2: Facilities Should Implement One or More of the Following Interventions/Strategies in All Blood Donors or in Subgroups of At-Risk Blood Donors (described below).

If blood collection establishments choose to implement one of the following interventions/strategies in only a single select donor subgroup, it is recommended that they begin with young donors (described below). However, it is anticipated that blood centers will progressively extend actions to all other at-risk groups.

- Subgroups at Risk

The following donors are at risk for developing iron deficiency and/or at increased risk for adverse effects of iron deficiency. In the absence of efforts to prevent iron deficiency in all donors, the following donors should be the target of interventions/strategies to mitigate iron deficiency:

- *Young donors.*

Adolescent donors remain a group at heightened risk of iron deficiency due to baseline iron requirements, growth, and dietary factors. At a minimum, this applies to donors 16-18 years of age. Evidence indicates that iron depletion may occur earlier and more frequently in this group, even at low donation intensity. It could also include donors through their early 20s, because substantive, measurable brain maturation and cognitive development can continue in that age group.³⁵⁻³⁷

It is therefore recommended that:

- Blood collection establishments adopt enhanced iron-protection strategies for adolescent donors as a default rather than an optional measure.
- Ferritin assessment should be considered before repeat donation in this age group.
- Donation frequency limits alone may be insufficient unless combined with iron supplementation or ferritin-guided policies.

- *Premenopausal females.*

- *Frequent donors, defined as:*



Advancing Transfusion and
Cellular Therapies Worldwide

- Males donating three or more times within a 12-month period.
 - Females donating two or more times within a 12-month period.
 - *Donors with hemoglobin values near the minimum for eligibility (eg, males with hemoglobin levels between 13.0 and 13.5 g/dL and females with hemoglobin levels between 12.5 and 13.0 g/dL).³⁸*
- Interventions/Strategies
When selecting among interventions, blood collection establishments should recognize that hemoglobin-only screening does not prevent iron deficiency and that lengthening donation intervals alone is unlikely to restore iron balance in most donors. Where operationally feasible, ferritin-guided strategies provide a targeted means of identifying donors at greatest risk and tailoring interventions accordingly.

These may include one or more of the following actions:

- Ferritin-Guided Donor Management: Operational Guidance
Ferritin testing provides a targeted approach that identifies donors who are demonstrably iron-depleted and allows personalized recommendations. Experience from large blood services highlights the importance of clear operational rules.

Recommended principles include:

- Ferritin testing in donor iron-management programs is typically performed on retained or stored plasma samples collected at the time of donation, thereby linking iron assessment directly to the donation encounter and ensuring standardized, contemporaneous evaluation of iron status (Sweegers et al, 2020).³⁹
- Donors with low ferritin should receive clear, written recommendations regarding iron supplementation and/or donation deferral.
- Deferral duration and supplementation advice should be linked to the degree of ferritin reduction, rather than using a single threshold.
- Re-entry criteria following low ferritin deferral should be defined prospectively to promote consistency and donor confidence.



Advancing Transfusion and
Cellular Therapies Worldwide

Clear communication is essential to prevent donor anxiety and to support informed donor participation.

○ *Donor iron supplementation.*

Low-dose oral iron supplementation⁴⁰ remains effective in replacing iron lost during donation. More recent physiological evidence indicates that iron absorption is influenced by hepcidin dynamics, suggesting that:

- Lower doses may achieve similar repletion with improved tolerability.
- Non-daily dosing schedules may be appropriate for donors who experience gastrointestinal intolerance.

Blood collection establishments may consider incorporating flexible supplementation strategies while maintaining the principle that iron replacement is necessary to restore iron balance following donation.

Blood collection establishments may choose to dispense iron supplements or multivitamins with iron directly to donors or provide vouchers exchangeable for iron to facilitate their obtaining iron supplements. Data from the HEIRS and STRIDE studies can be used to determine the minimal iron dose and duration of administration. As shown in HEIRS, a 38-mg supplement resulted in the majority of iron being replaced within the first 8 weeks; in the STRIDE study, 19- and 38-mg doses were equally effective in replacing iron lost from blood donation. Therefore, to replace the iron lost in a WB donation, a minimum of 18 mg of elemental iron [as a multivitamin with iron (18 mg being the most commonly available dosage) or an iron supplement] is recommended daily for 60 days.

Based on the amount of red cells removed at donation, it seems logical that 2-unit Red Blood Cell (RBC) donors should take iron for twice as long as single-unit donors. However, iron absorption is quite dependent on donor iron status; ie, donors with a greater degree of iron deficiency will absorb more iron. Therefore, 2-unit RBC donors may also get the maximal benefit from supplemental iron in the first 8 weeks after donation when they have their lowest ferritin levels. Currently, in the absence of data that address this issue, blood centers should provide 2-unit RBC donors with supplements or vouchers for a minimum of 18 mg of elemental iron daily for either 60 or 120 days following donation, at the discretion of the facility's medical director.



Advancing Transfusion and
Cellular Therapies Worldwide

Blood centers should define the number of apheresis procedures (platelets and plasma) that result in red cell losses (including incomplete procedures and blood samples) that approximate the red cell loss in one WB donation. Based upon kit residuals and testing tube losses, this is likely to be four to five plateletpheresis or plasmapheresis donations.^{41, 42} Donors reaching red cell losses equivalent to those of frequent blood donors should be informed about iron loss and provided with supplements or vouchers for a minimum of 18 mg of elemental iron daily for 60 days; this same regimen should be repeated at intervals approximating each subsequent equivalent WB unit loss.

With the introduction of iron supplementation programs, it is expected that blood collection establishments will experience a reduction in low hemoglobin deferrals. A 2014 Cochrane meta-analysis of four studies of iron supplementation in blood donors describes a 66% reduction in deferrals with iron use (95% confidence interval 45% - 79%).⁴³

- *Lengthening the interdonational interval and/or decreasing the number of donations per year.*

Blood collection establishments may choose to increase interdonation intervals and/or decrease the maximal permissible number of donations per year. There is clear evidence from observational studies that the number of donations in the previous 1-2 years is the foremost determinant of iron deficiency in blood donors.² Furthermore, it is well-documented that a 56-day interdonation interval, without iron supplementation, produces a negative iron balance.⁴⁴ Even countries with an interdonation interval greater than the 56 days permitted in North America have reported high rates of hemoglobin deferral due to iron deficiency in the absence of iron supplementation.⁴⁵ In the HEIRS study, two-thirds of subjects receiving no iron supplementation had not recovered lost iron by 24 weeks after donation.²⁵ Based upon these and other data,⁴⁴ it is increasingly clear that lengthening the interdonation interval is highly unlikely to be fully effective as a solitary intervention to mitigate donor iron deficiency unless the interval is lengthened to ≥ 26 weeks. In other words, if no other interventions in addition to extended interdonation interval or limitations on donation frequency are utilized as a strategy to protect blood donors from iron deficiency, then limiting WB donations to two times in a 12-month period or one 2-unit RBC donation in a 12-month period appears to be the maximal collection schedule that can be employed. Because of higher documented rates of iron deficiency in young donors, it is recommended that in the absence of any other intervention for such donors (ages 16-18) of both genders, blood collection



Advancing Transfusion and
Cellular Therapies Worldwide

establishments should limit collections to the red cell loss from one WB donation in a 12-month period. Collections from premenopausal women may similarly be limited to one donation in a 12-month period at the discretion of the facility's medical director.

Limiting the number of donations per year to this extent is likely to have a significant impact on the adequacy of the blood supply in the absence of a significantly expanded donor base.^{46, 47} Therefore, blood collection establishments may choose to combine a lesser limitation on donation intervals or frequency with other strategies listed in this Association Bulletin.

- *Donor ferritin testing as a basis for advising donors about further actions.*

Blood collection establishments may choose to test ferritin levels for all donors in defined or donor subgroups to guide personalized recommendations for iron supplementation before the development of anemia. This allows for a targeted approach to donors demonstrated to be iron deficient rather than targeting donors presumed to be iron deficient by a demographic characteristic or prior donation activity. Experience from large-scale implementation programs indicates that ferritin-guided management is most effective when deferral duration, supplementation advice, and criteria for return to donation are prospectively defined and linked to the degree of ferritin reduction. Based on findings from the STRIDE study, it appears that donors may be highly motivated to follow blood center iron supplementation recommendations or self-extend interdonation intervals when their ferritin value documents iron deficiency. Canadian data outside the more carefully controlled STRIDE environment highlight the need for especially clear communication.⁴⁸

Ferritin testing should be performed on a sample collected at the time of blood donation. A low ferritin result would apply to management of the donor before any subsequent donation. Clear recommendations for iron supplementation and/or for interdonation interval extension should be provided in writing to donors along with their ferritin test results. Repeat ferritin testing should be determined by prior test results and operational capabilities of the blood collection establishment.

Adolescents and Young Donors

Adolescent donors (ages 16–18) represent a group at increased risk of iron deficiency due to ongoing growth, baseline iron requirements, and lower iron reserves. Evidence indicates that iron depletion may occur earlier and with fewer donations in this population.⁵⁰ Blood collection establishments should therefore consider enhanced iron-protection strategies for adolescent donors



Advancing Transfusion and
Cellular Therapies Worldwide

as a default approach. Such strategies may include ferritin assessment prior to repeat donation, provision of iron supplementation, or stricter limits on donation frequency unless iron status is demonstrated to be adequate.

Recommendation #3: Blood Collection Establishments Should Consider Postimplementation Monitoring

Blood collection establishments are encouraged to collect, track, tabulate, and publish data on any interventions they conduct. AABB also encourages blood collection establishments to design their interventions in such a way that endpoints are clear and the derived data can be readily compared to interventions implemented at other facilities.

Post-implementation monitoring should extend beyond laboratory indices to include operational and donor-centered outcomes.

Suggested metrics include:

- Prevalence of low ferritin over time.
- Rates of low-hemoglobin deferral.
- Donor return and retention.
- Where feasible, donor-reported symptoms related to iron deficiency.

Such data allow continuous refinement of donor protection strategies and facilitate comparison across programs.

Blood collection establishments are encouraged to collect, track, tabulate, and publish data on any interventions they conduct. AABB also encourages blood collection establishments to design their interventions in such a way that endpoints are clear and the derived data can be readily compared to interventions implemented at other facilities.



Advancing Transfusion and
Cellular Therapies Worldwide

References

1. Noone, S, ed. Standards for blood banks and transfusion services. 35th ed. Bethesda, MD: AABB, 2026.
2. Cable RG, Glynn SA, Kiss JE, et al for the NHLBI Retrovirus Epidemiology Donor Study-II (REDS-II). Iron deficiency in blood donors: Analysis of enrollment data from the REDS-II Donor Iron Status Evaluation (RISE) study. *Transfusion* 2011;51:511-22.
3. Cable RG, Glynn SA, Kiss JE, et al for the NHLBI Retrovirus Epidemiology Donor Study-II (REDS-II). Iron deficiency in blood donors: The REDS-II Donor Iron Status Evaluation (RISE) study. *Transfusion* 2012;52:702-11.
4. Pratt JJ, Khan KS. Non-anaemic iron deficiency – a disease looking for recognition of diagnosis: A systematic review. *Eur J Haematol* 2015;96:618-28.
5. Sohrabi F. Tip of the iceberg: Extra-haematological consequences of early iron deficiency. *J Glob Health* 2015;5(2):020304.
6. Bryant BJ, Yau YY, Arceo SM, et al. Ascertainment of iron deficiency and depletion in blood donors through screening questions for pica and restless legs syndrome. *Transfusion* 2013;53:1637-44.
7. Spencer BR, Kleinman S, Wright DJ, et al. Restless legs syndrome, pica, and iron status in blood donors. *Transfusion* 2013;53:1645-52.
8. Silber MH, Richardson JW. Multiple blood donations associated with iron deficiency in patients with restless legs syndrome. *Mayo Clin Proc* 2003;78:52-4.
9. Krayenbuehl PA, Battegay E, Breymann C, et al. Intravenous iron for the treatment of fatigue in nonanemic, premenopausal women with low serum ferritin concentration. *Blood* 2011;118:3222-7.
10. Vaucher P, Druais PL, Waldvogel S, et al. Effect of iron supplementation on fatigue in nonanemic menstruating women with low ferritin: A randomized controlled trial. *CMAJ* 2012;184:1247-54.



Advancing Transfusion and
Cellular Therapies Worldwide

11. Pittori C, Buser A, Gasser UE, et al. A pilot iron substitution programme in female blood donors with iron deficiency without anemia. *Vox Sang* 2011;100:303-11.
12. Waldvogel S, Pedrazzini B, Vaucher P, et al. Clinical evaluation of iron treatment efficiency among non-anemic but iron-deficient female blood donors: A randomized controlled trial. *BMC Med* 2012;10:8.
13. Rigas AS, Pedersen OB, Sørensen CJ, et al. No association between iron status and self-reported health-related quality of life in 16,375 Danish blood donors: Results from the Danish Blood Donor Study. *Transfusion* 2015;55:1752-6.
14. Breymann C. Iron deficiency anemia in pregnancy. *Semin Hematol* 2015;52(4):339-47.
15. McCann JC, Ames BN. An overview of evidence for a causal relation between iron deficiency during development and deficits in cognitive or behavioral function. *Am J Clin Nutr* 2007;85:931-45.
16. Ribot B, Aranda N, Viteri F, et al. Depleted iron stores without anaemia early in pregnancy carries increased risk of lower birthweight even when supplemented daily with moderate iron. *Hum Reprod* 2012;27:1260-66.
17. Jáuregui-Lobera I. Iron deficiency and cognitive functions. *Neuropsych Dis Treat* 2014;10:2087-95.
18. Falkingham M, Abdelhamid A, Curtis P, et al. The effects of oral iron supplementation on cognition in older children and adults: A systematic review and meta-analysis. *Nutr J* 2010;9:4.
19. Murray-Kolb LE, Beard JL. Iron treatment normalizes cognitive functioning in young women. *Am J Clin Nutr* 2007;85:778-87.
20. Stugiewicz M, Tkaczyszyn M, Kasztura M, et al. The influence of iron deficiency on the functioning of skeletal muscles: Experimental evidence and clinical implications. *Eur J Heart Fail* 2016;18:762-73.



Advancing Transfusion and
Cellular Therapies Worldwide

21. Saraon T, Katz SD. Targeting iron deficiency anemia in heart failure. *Prog Cardiovasc Dis* 2016;58:407-15.
22. von Haehling S, Jankowska WA, van Veldhuisen DJ, et al. Iron deficiency and cardiovascular disease. *Nat Rev Cardiol* 2015;12:659-69.
23. Bruner A, Joffe A, Duggan A, et al. Randomised study of cognitive effects of iron supplementation in non-anaemic iron-deficient adolescent girls. *Lancet* 1996;348:992-6.
24. Domen RE. Drugs and donors: Ethical considerations in the use of hematopoietic enhancing agents in volunteer whole blood and blood component donors. In: Domen RE, ed. *Ethical issues in transfusion medicine and cellular therapies*. Bethesda, MD: AABB Press, 2015:99-108.
25. Kiss JE, Brambilla D, Glynn SA, et al for the NHLBI REDS-III. Oral iron supplementation after blood donation: A randomized clinical trial. *JAMA* 2015;313(6):575-83.
26. Mast AE, Bialkowski W, Bryant BJ, et al. A randomized, blinded, placebo-controlled trial of education and iron supplementation for mitigation of iron deficiency in regular blood donors. *Transfusion* 2016;56(6 Pt 2):1588-97.
27. Food and Drug Administration. Blood Product Advisory Committee meeting webcast, November 17, 2017. [Available at: <https://collaboration.fda.gov/p9854v7bgrh/?launcher=false&fcsContent=true&pbMode=normal>.]
28. Cable RG, Brambilla D, Glynn SA, et al for the NHLBI REDS-III. Effect of iron supplementation on iron stores and total body iron after whole blood donation. *Transfusion* 2016;56:2005-12.
29. Vinkenoog M, van den Hurk K, van Kraaij M, van Leeuwen M, Janssen MP. First results of a ferritin-based blood donor deferral policy in the Netherlands. *Transfusion*. 2020 Aug;60(8):1785-1792. doi: 10.1111/trf.15906. Epub 2020 Jun 13. PMID: 32533600; PMCID: PMC7496980.
30. Karregat JHM, Meulenbeld A, Quee FA, Novotny VMJ, Swinkels DW, van den Born BH, Zaaijer HL, Twisk JWR, van den Hurk K. Ferritin-guided iron supplementation as an



Advancing Transfusion and
Cellular Therapies Worldwide

- alternative or complement to prolonged blood donation intervals (FORTE): a double-blind, randomised, controlled trial. *Lancet Haematol*. 2025 Sep;12(9):e694-e704. doi: 10.1016/S2352-3026(25)00167-X. Epub 2025 Aug 14. PMID: 40819648.
31. Karregat J, Sweegers MG, Quee FA, Weekamp HH, Swinkels DW, Novotny VMJ, Zaaijer HL, van den Hurk K. Ferritin-guided iron supplementation in whole blood donors: optimal dosage, donor response, return and efficacy (FORTE)-a randomised controlled trial protocol. *BMJ Open*. 2022 Mar 9;12(3):e056316. doi: 10.1136/bmjopen-2021-056316. PMID: 35264362; PMCID: PMC8915278.
 32. Drechsler LØ, Boldsen JK, Hindhede L, Aagaard B, Harritshøj LH, Mikkelsen C, Brodersen T, Brøns N, Schwinn M, Hjalgrim H, Rostgaard K, Topholm Bruun M, Ostrowski SR, Pedersen OB, Mikkelsen S, Erikstrup C. The effect of ferritin-guided iron supplementation among Danish female first-time blood donors. *Transfusion*. 2023 Sep;63(9):1710-1718. doi: 10.1111/trf.17484. Epub 2023 Jul 14. PMID: 37452554
 33. van den Hurk K, Arvas M, Roberts DJ, Castrén J, Erikstrup C. Whole Blood Donor Iron Management Across Europe: Experiences and Challenges in Four Blood Establishments. *Transfus Med Rev*. 2024 Oct;38(4):150860. doi: 10.1016/j.tmr.2024.150860. Epub 2024 Sep 18. PMID: 39369584.
 34. von Siebenthal HK, Gessler S, Vallelia F, Steinwendner J, Kuenzi UM, Moretti D, Zimmermann MB, Stoffel NU. Alternate day versus consecutive day oral iron supplementation in iron-depleted women: a randomized double-blind placebo-controlled study. *EClinicalMedicine*. 2023 Nov 3;65:102286. doi: 10.1016/j.eclinm.2023.102286. PMID: 38021373; PMCID: PMC10659995.
 35. Luna B, Marek S, Larsen B, et al. An integrative model of the maturation of cognitive control. *Annu Rev Neurosci* 2015;38:151-70.
 36. Mills KL, Goddings AL, Clasen AS, et al. The developmental mismatch in structural brain maturation during adolescence. *Dev Neurosci* 2014;36:147-60.
 37. Casey BJ, Jones RM, Somerville LH. Braking and accelerating of the adolescent brain. *J Res Adolesc* 2011;21:21-33.



Advancing Transfusion and
Cellular Therapies Worldwide

38. Bravo MD, Custer B, Tomasulo PA, et al. Age and gender relationships observed in routine ferritin testing (abstract). *Transfusion* 2014;54(2S):116A-7A.
39. Sweegers MG, Zalpur S, Quee FA, Huis In 't Veld EMJ, Prinsze FJ, Hoogendijk EO, Twisk JWR, van Weert AWM, de Kort WLAM, van den Hurk K. Ferritin measurement IN Donors-Effectiveness of iron Monitoring to diminish iron deficiency and low haemoglobin in whole blood donors (FIND'EM): study protocol for a stepped wedge cluster randomised trial. *Trials*. 2020 Oct 1;21(1):823. doi: 10.1186/s13063-020-04648-w. PMID: 32998766; PMCID: PMC7528293.
40. Kiss JE, Brambilla D, Glynn SA, Mast AE, Spencer BR, Stone M, Kleinman SH, Cable RG; National Heart, Lung, and Blood Institute (NHLBI) Recipient Epidemiology and Donor Evaluation Study-III (REDS-III). Oral iron supplementation after blood donation: a randomized clinical trial. *JAMA*. 2015 Feb 10;313(6):575-83. doi: 10.1001/jama.2015.119. PMID: 25668261; PMCID: PMC5094173.
41. Li H, Condon F, Kessler D, et al. Evidence of relative iron deficiency in platelet- and plasma-pheresis donors correlates with donation frequency. *J Clin Apher* 2016;31:551-8.
42. Page EA, Coppock JE, Harrison JF. Study of iron stores in regular plateletpheresis donors. *Transfus Med* 2010;20:22-9.
43. Smith GA, Fisher SA, Doree C. et al. Oral or parenteral iron supplementation to reduce deferral, iron deficiency and/or anaemia in blood donors. *Cochrane Database Syst Rev* 2014;(7):CD009532.
44. Schotten N, Pasker-de Jong PCM, Moretti D, et al. The donation interval of 56 days requires extension to 180 days for whole blood donors to recover from changes in iron metabolism. *Blood* 2016;128:2185-8.
45. Goldman M, Magnussen K, Gorlin J, et al. International forum regarding practices related to donor haemoglobin and iron. *Vox Sang* 2016;111:449-55.
46. Spencer BR, Johnson B, Wright DJ, et al for the REDS-II RISE Analysis Group. Potential impact on blood availability and donor iron status of changes to donor hemoglobin cutoff and interdonation intervals. *Transfusion* 2016;56:1994-2004.



Advancing Transfusion and
Cellular Therapies Worldwide

47. Sayers M, Centilli J. Contemplating the effect on blood availability if the interdonation interval of 56 days is prolonged. *Transfusion* 2013;53:1132-6.
48. Goldman M, Uzicanin S, Scalia J, et al. Impact of informing donors of low ferritin results. *Transfusion* 2016;56:2193-8.
50. Patel EU, White JL, Bloch EM, Grabowski MK, Gehrie EA, Lokhandwala PM, Brunker PAR, Goel R, Shaz BH, Ness PM, Tobian AAR. Association of blood donation with iron deficiency among adolescent and adult females in the United States: a nationally representative study. *Transfusion*. 2019 May;59(5):1723-1733. doi: 10.1111/trf.15179. Epub 2019 Feb 18. PMID: 30779173; PMCID: PMC6791124.



Advancing Transfusion and
Cellular Therapies Worldwide

ADDENDUM

Suggested Donor Iron Management Algorithm

It is recommended that blood services set up their own donor iron management algorithms based on the following example.

Step 1: Identify donors requiring iron-protection strategies

Iron-protection strategies should be applied to donors in any of the following categories:

- **Adolescents and young donors (minimum ages 16–18).**
- **Premenopausal females.**
- **Frequent donors (≥ 3 whole blood donations per year in males; ≥ 2 in females, or equivalent red cell loss).**
- **Donors with hemoglobin values near minimum eligibility thresholds.**
- **Apheresis donors with cumulative red cell losses approaching whole blood equivalence.**

Step 2: Assess iron status or apply default protection

Blood collection establishments may:

- **Measure ferritin at the time of donation to guide donor-specific management; or**
- **Apply default iron supplementation and/or donation-interval modification in defined high-risk groups where ferritin testing is not performed.**

Step 3: Interpret ferritin and determine action (example framework)

While specific thresholds may vary, a graded approach is recommended:

- **Markedly low ferritin:**
Donation deferral for a defined interval plus iron supplementation.
- **Low ferritin:**
Shorter deferral and/or iron supplementation with defined re-entry criteria.
- **Low-normal ferritin:**
Donation may proceed with recommendation for post-donation iron supplementation and/or extended interval.



Advancing Transfusion and
Cellular Therapies Worldwide

- **Adequate ferritin:**
Routine donation permitted with standard donor education.

Step 4: Iron supplementation guidance

- **Low-dose oral iron is effective in replacing iron lost through donation.**
- **Supplementation duration should reflect donation intensity and iron status.**
- **Alternative dosing schedules may be considered for donors with intolerance, provided adequate iron repletion is achieved.**

Step 5: Define re-entry criteria

Donors deferred for low ferritin should receive:

- **Written guidance on supplementation and deferral duration.**
- **Clear criteria for return to donation, based on time elapsed, ferritin recovery, or both.**

Step 6: Monitor program outcomes

Blood collection establishments should monitor:

- **Prevalence of low ferritin and iron deficiency.**
- **Rates of hemoglobin deferral.**
- **Donor return and retention.**
- **Where feasible, donor-reported symptoms related to iron deficiency.**